


CARE4 MODULE NAME: PCS

Pausing/Resuming IV Infusions (Including Heparin)

For complete workflow for eMAR IV infusions including heparin, please review eMAR IV Infusions Tip Sheet on the CARE4 App or website.


Pausing IV Infusions

Enter Infusion Rate of "0". Increase/Decrease field updates to "Paused". Select Save and this will update on eMAR as "paused".



Once you enter "0" in the infusion rate field (required field) it auto pauses the IV/medication/heparin.


sodium chloride 0.9 % 1,000 ml @ 100 mls/hr IV .Q10H SCH		
Current Rate: 100 mls/hr		
U00072572 M Q Inf FULL		
Infusion/Titration		
IV site		
IV Site	Right Antecubital	
Intake		
Intake		36.667
Cumulative In...		36.667
Cumulative In...		36.667
Container Vol...	1,000	963.333
Waste Amount	0	0
Dosing		
Dose Rate		
Infusion Rate ...	100	0
Cumulative Dose	Not Applicable	Not Applicable
Increase/Decr...	Started	Paused
Stop/Elapsed Time		
Elapsed Time ...	0m	22m
Stop/Total Do...		



Restarting Paused IV Infusions

Enter Infusion Rate at ordered rate. Increase/Decrease field updates to Resumed. Other fields will auto populate and continue with previous amounts. Select Save and Save

Document Assess			Document Inf/Titr		
		Thu 23 Sep 20:34 AA			Thu 23 Sep 20:59 AA
sodium chloride 0.9 % 1,000 ml @ 100 mls/hr IV .Q10H SCH					
Current Rate: 100 mls/hr					
U00072572 M Q Inf FULL					
Infusion/Titration					
IV site					
IV Site	Right Antecubital				
Intake					
Intake		36.667		0	
Cumulative In...		36.667		36.667	
Cumulative In...		36.667		36.667	
Container Vol...	1,000	963.333		963.333	
Waste Amount	0	0		0	
Dosing					
Dose Rate					
Infusion Rate ...	100	0		100	
Cumulative Dose	Not Applicable		Not Applicable		Not Applicable
Increase/Decr...	Started		Paused		Resumed
Stop/Elapsed Time					
Elapsed Time ...	0m	22m		22m	
Stop/Total Do...					



Ending/Stopping IV Infusions – No Wastage, bag is empty

- Enter “0” in “Container Volume” field of Intake section
- Dosing section, “Increase/Decrease” field updates to “Infused”
- Other field’s auto populate. Save.

Note – when there is less than 10% bag volume remaining, scanning the new IV bag will automatically end the previous infusion and start the new infusion. If there is more than 10% of the bag volume remaining, the user must end the previous bag following Ending/Stopping IV Infusions procedure.

Ending/Stopping IV Infusions when the bag is not empty – Documenting Wastage

Enter Infusion Rate of “0”. Copy the remaining Container Volume amount into the Waste Amount field. The IV will now display as infused on the eMAR.